

The Leading Coefficient of Certain Kazhdan-Lusztig Polynomials of the Permutation Group $\mathfrak{S}_n$

Nanhua Xi

ABSTRACT. In this paper we show that the leading coefficient $\mu(y, w)$ of certain Kazhdan-Lusztig polynomials $P_{y,w}$ of the permutation group $\mathfrak{S}_n$ of $1, 2, \dots, n$ are not greater than 1. More precisely, we show that the leading coefficients $\mu(y, w)$ are not greater than 1 whenever $a(y) < a(w)$, where $a : \mathfrak{S}_n \rightarrow \mathbf{N}$ is the function defined in [L1]. See 1.5 for a simple interpretation of the function a .

The Kazhdan-Lusztig polynomials of a Coxeter group (W, S) play a central role in Kazhdan-Lusztig theory. From the seminal work [KL] one sees that the leading coefficient $\mu(y, w)$ of some Kazhdan-Lusztig polynomials $P_{y,w}$ are of great importance. It is certainly interesting to understand the leading coefficients. We have two natural questions, (a) determine the leading coefficients $\mu(y, w)$, (b) for a given element w in the Coxeter group, determine all y such that $\mu(y, w)$ or $\mu(w, y)$ is non-zero. By now, for both questions we do not have much understanding except for some special cases, see for example [L4].

In this paper we are concerned with question (a) for the permutation group $\mathfrak{S}_n$ of $1, 2, \dots, n$. We shall show that for $y, w \in \mathfrak{S}_n$, if $y \leq w$ (Bruhat order) and $a(y) < a(w)$, then the leading coefficient $\mu(y, w)$ is less than or equal to 1, see Theorem 1.4, here the function $a : \mathfrak{S}_n \rightarrow \mathbf{N}$ is defined by Lusztig in [L1], see 1.5 for a simple interpretation of the function for $\mathfrak{S}_n$. When $a(y) \geq a(w)$, McLarnan and Warrington have showed that $\mu(y, w)$ can be greater than 1, see [MW]. The work [MW] gives a negative answer to the (0,1)-conjecture in [GM, p.52].

In this paper we also give some discussion for the set of all $\mu(y, w)$, $y, w \in \mathfrak{S}_n$, see section 1.2 and 3.2.

1. Preliminaries

In this section we collect some terms and basic facts, we refer to [KL] for more details.

1.1. Let (W, S) (S the set of simple reflections) be a Coxeter group. We shall denote the length function of W by l and use $\leq$ for the Bruhat order on W .

Let H be the Hecke algebra of (W, S) over $\mathcal{A} = \mathbf{Z}[q^{\frac{1}{2}}, q^{-\frac{1}{2}}]$ (q an indeterminate) with parameter q . Let $\{T_w\}_{w \in W}$ be its standard basis. There is a unique ring homomorphism $\bar{\cdot} : H \rightarrow H$ such that $\overline{T_w} = T_{w^{-1}}$ and $\bar{q}^{\frac{1}{2}} = q^{-\frac{1}{2}}$.

In [KL1], Kazhdan and Lusztig proved that for each w in W , there exists a unique element C_w in the Hecke algebra H such that (1) $C_w = \bar{C}_w$, (2) $C_w = q^{-\frac{l(w)}{2}} \sum_{y \leq w} P_{y,w} T_y$, where $P_{y,w}$ is a polynomial in q of degree less than or equal to $\frac{1}{2}(l(w) - l(y) - 1)$ for $y < w$ and $P_{w,w} = 1$.

The basis $\{C_w\}_{w \in W}$ of H is the Kazhdan-Lusztig basis. The polynomials $P_{y,w}$ are the Kazhdan-Lusztig polynomials. If $y < w$, we have

$$P_{y,w} = \mu(y, w)q^{\frac{1}{2}(l(w)-l(y)-1)} + \text{lower degree terms.}$$

We write $y \prec w$ if $\mu(y, w) \neq 0$.

1.2. Assume that (W, S) is an affine Weyl group or a Weyl group. The following formula is due to Springer (see [Sp]),

$$(a) \quad \mu(y, x) = \sum_{d \in \mathcal{D}_0} \delta_{y^{-1}, x, d} + \sum_{f \in \mathcal{D}_1} \gamma_{y^{-1}, x, f} \pi(f) = \sum_{d \in \mathcal{D}_0} \delta_{y, x^{-1}, d} + \sum_{f \in \mathcal{D}_1} \gamma_{y, x^{-1}, f} \pi(f).$$

We need explain the notations. Write

$$C_x C_y = \sum_{z \in W} h_{x, y, z} C_z, \quad h_{x, y, z} \in \mathcal{A} = \mathbf{Z}[q^{\frac{1}{2}}, q^{-\frac{1}{2}}].$$

Following Lusztig ([L1]), we define

$$a(z) = \min\{i \in \mathbf{N} \mid q^{-\frac{i}{2}} h_{x, y, z} \in \mathbf{Z}[q^{-\frac{1}{2}}] \text{ for all } x, y \in W\}.$$

If for any i , $q^{-\frac{i}{2}} h_{x, y, z} \notin \mathbf{Z}[q^{-\frac{1}{2}}]$ for some $x, y \in W$, we set $a(z) = \infty$. It is not clear that whether there exists a Coxeter group (W, S) such that $a(z) = \infty$ for some $z \in W$.

From now on, we assume that the function $a : W \rightarrow \mathbf{N}$ is bounded and (W, S) is crystallographic. In this case, the function a is constant on two-sided cells of W (see [L1]). Obviously, when W is finite, the function a is bounded. Lusztig showed that the function a is bounded for all affine Weyl groups (see [L1]). Following Lusztig and Springer, we define $\delta_{x, y, z}$ and $\gamma_{x, y, z}$ by the following formula,

$$h_{x, y, z} = \gamma_{x, y, z} q^{\frac{a(z)}{2}} + \delta_{x, y, z} q^{\frac{a(z)-1}{2}} + \text{lower degree terms.}$$

Springer showed that $l(z) \geq a(z)$ (see [L2]). Let $\delta(z)$ be the degree of $P_{e, z}$, where e is the neutral element of W . Then actually one has $l(z) - a(z) - 2\delta(z) \geq 0$ (see [L2]). Set

$$\mathcal{D}_i = \{z \in W \mid l(z) - a(z) - 2\delta(z) = i\}.$$

The number $\pi(z)$ is defined by $P_{e, z} = \pi(z)q^{\delta(z)} + \text{lower degree terms}$.

The elements of $\mathcal{D}_0$ are involutions, called distinguished involutions of (W, S) (see [L2]). Moreover, in a Weyl group or an affine Weyl group, each left cell (resp. right cell) contains exactly one element of $\mathcal{D}_0$, see [L2]. When W is a Weyl group, Lusztig has showed that an element z is in $\mathcal{D}_1$ if and only if there exists some d in $\mathcal{D}_0$ such that $z \underset{LR}{\sim} d$ and $\mu(z, d) \neq 0$ or $\mu(d, z) \neq 0$, see [L3]. For $\mathfrak{S}_n$, all involutions are distinguished.

Example: Let S' be a subset of S such that the subgroup W' of W generated by S' is finite. Then the longest element of W' is in $\mathcal{D}_0$.

1.3. We refer to [KL] for the definition of the preorders $\leq_L, \leq_R, \leq_{LR}$ and of the equivalence relations $\underset{L}{\sim}, \underset{R}{\sim}, \underset{LR}{\sim}$ on W . The corresponding equivalence classes are called left cells, right cells, two-sided cells of W , respectively. The preorder $\leq_L$ (resp. $\leq_R, \leq_{LR}$) induces a partial order on the set of left (resp. right; two-sided) cells of W , denoted again by $\leq_L$ (resp. $\leq_R, \leq_{LR}$). For a Weyl group or an affine Weyl group, Springer showed the following results (a) and (b) (see [Sp])

- (a) Assume that $\mu(y, w)$ or $\mu(w, y)$ is nonzero, then $y \leq_L w$ and $y \leq_R w$ if $a(y) < a(w)$, and $y \sim_L w$ or $y \sim_R w$ if $a(y) = a(w)$.
- (b) If $\delta_{x,y,z} \neq 0$, then $z \sim_L y$ or $z \sim_R x$. (Note that $h_{x,y,z} \neq 0$ implies that $a(z) \geq a(x)$ and $a(z) \geq a(y)$, see [L1].)
- For $w \in W$, set $L(w) = \{s \in S \mid sw \leq w\}$, $R(w) = \{s \in S \mid ws \leq w\}$. Then we have (see [KL])
- (c) $R(w) \subseteq R(y)$ if $y \leq_L w$. In particular, $R(w) = R(y)$ if $y \sim_L w$;
- (d) $L(w) \subseteq L(y)$ if $y \leq_R w$. In particular, $L(w) = L(y)$ if $y \sim_R w$.

We shall need the star operation introduced in [KL]. Let s and t be in S such that st has order 3, i.e. $sts = tst$. Define

$$D_L(s, t) = \{w \in W \mid L(w) \cap \{s, t\} \text{ has exactly one element}\},$$

$$D_R(s, t) = \{w \in W \mid R(w) \cap \{s, t\} \text{ has exactly one element}\}.$$

If w is in $D_L(s, t)$, then $\{sw, tw\}$ contains exactly one element in $D_L(s, t)$, denoted by $*w$, here $*$ = $\{s, t\}$. The map: $\varphi : D_L(s, t) \rightarrow D_L(s, t)$, $w \rightarrow *w$, is an involution and is called a left star operation. Similarly if $w \in D_R(s, t)$ we can define the right star operation $\psi : D_R(s, t) \rightarrow D_R(s, t)$, $w \rightarrow w^* = \{ws, wt\} \cap D_R(s, t)$, where $*$ = $\{s, t\}$. The following result is proved in [KL].

- (e) Let s and t be in S such that st has order 3. Assume that $y \prec w$. If both y^* and w^* (resp. $*y$ and $*w$) are well defined, then either (1) $y^* \prec w^*$ and $\mu(y, w) = \mu(y^*, w^*)$, or (2) $w^* \prec y^*$ and $\mu(y, w) = \mu(w^*, y^*) = 1$ (resp. either (1) $*y \prec *w$ and $\mu(y, w) = \mu(*y, *w)$, or (2) $*w \prec *y$ and $\mu(y, w) = \mu(*w, *y) = 1$).

The main result of this paper is the following.

1.4. Theorem. Let $y, w \in \mathfrak{S}_n$. If $y < w$ and $a(y) < a(w)$, then the leading coefficient $\mu(y, w)$ is less than or equal to 1, here the function $a : \mathfrak{S}_n \rightarrow \mathbf{N}$ is defined in 1.2.

Remark. This result is also true for an affine Weyl group of type $\tilde{A}_{n-1}$.

1.5. For $W = \mathfrak{S}_n$, there is a nice description for cells of W , called Robinson-Schensted rule. We recall the rule since it also provide a simple way to compute the function $a : \mathfrak{S}_n \rightarrow \mathbf{N}$ defined in subsection 1.2. As usual, the set S consists of transpositions $s_i = (i, i+1)$, $i = 1, 2, \dots, n-1$.

A partition λ of n is visualized as a Young diagram F_λ with n boxes such that the lengths of the columns of the diagram are given by the components of λ . A Young tableau of shape λ is a labelling of the boxes of F_λ with integers $1, 2, \dots, n$. The tableau is called standard if the numbers labelling its boxes are increasing in the columns from the top to the bottom and are increasing in the row from left to right. The Robinson-Schensted rule gives a bijection $\theta : w \rightarrow (P(w), Q(w))$ from the permutation group $\mathfrak{S}_n$ to the set of pairs of standard Young tableau on $\{1, 2, \dots, n\}$. Let $w(i) = j_i$ for $i = 1, 2, \dots, n$. The standard Young tableau $P(w)$ is defined as follows.

Let T_0 be the empty Young tableau. Assume that we have defined a Young tableau T_{i-1} which has totally $i-1$ boxes and the boxes are labelled with integers $j_1, j_2, \dots, j_{i-1}$. Now we define the Young tableau T_i . If j_i is greater than every numbers in the first row of T_{i-1} , then T_i is obtained from T_{i-1} by adding j_i to the end of the first row of T_{i-1} . If not, then let z be the smallest number of the first

row of T_{i-1} greater than j_i . Let T'_{i-1} be the Young tableau by removing the first row of T_{i-1} . Then T_i is the Young tableau obtained from T'_{i-1} by replacing z by j_i and by inserting z to the tableau T'_{i-1} as the same way. It is known that $P(w)$ is indeed a standard Young tableau. We set $Q(w) = P(w^{-1})$. Let y, w be in $\mathfrak{S}_n$, then we have (see [BV])

- (a) $y \underset{L}{\sim} w$ if and only if $Q(y) = Q(w)$,
- (b) $y \underset{R}{\sim} w$ if and only if $P(y) = P(w)$,
- (c) $y \underset{LR}{\sim} w$ if and only if $P(y)$ and $P(w)$ have the same shape.

Assume that $P(w)$ is a Young tableau of shape $\lambda = (\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_k)$ (a partition of n). we define $a(w)$ to be $\frac{1}{2} \sum_{i=1}^k \lambda_i(\lambda_i - 1)$. Then $a : \mathfrak{S}_n \rightarrow \mathbf{N}$ is exactly the a -function defined in 1.2.

Note that $y \underset{LR}{\leq} w$ if and only if $\lambda(y) \leq \lambda(w)$, where $\lambda(y)$ and $\lambda(w)$ are the partitions corresponding to $P(y)$ and $P(w)$ respectively. (We say $\lambda = (a_1, a_2, \dots) \leq \mu = (b_1, b_2, \dots)$ if $a_1 + \dots + a_k \leq b_1 + \dots + b_k$ for all $k = 1, 2, \dots$)

2. A description of the set of all $\mu(y, w)$ for $\mathfrak{S}_n$

In this section we give a description to the set of all $\mu(y, w)$, $y, w \in \mathfrak{S}_n$, in terms of $\delta_{x,y,z}$, $x, y, z \in \mathfrak{S}_n$. More precisely, we have

Proposition 2.1. For $\mathfrak{S}_n$, the set $\{\delta_{x,y,z} | x, y, z \in \mathfrak{S}_n\}$ is equal to the set $\{\mu(y, w) | y, w \in \mathfrak{S}_n\}$.

Proof: Clearly both sets contain 0. Assume that $\mu(y, w)$ is non zero. By Springer's formula 1.2(a), we have $\delta_{y^{-1},w,d} \neq 0$ for some $d \in \mathcal{D}_0$ or $\gamma_{y^{-1},w,u} \neq 0$ for some $u \in \mathcal{D}_1$. It is no harm to assume that $a(w) \geq a(y)$. If $a(w) > a(y)$, then $\gamma_{y^{-1},w,u} = 0$ for all u in $\mathfrak{S}_n$ (see [L1, 6.3(a)]), and $\delta_{y^{-1},w,d} \neq 0$ implies that d and w are in the same left cell, cf 1.2 (c) and [L1, 6.3(b)]. Therefore $\mu(y, w) = \delta_{y^{-1},w,d(w)}$ in this case. (We use $d(x)$ for the unique element in $\mathcal{D}_0$ which is in the left cell containing x .) If $a(w) = a(y)$, by 1.3 (a), then w and y are in the same left cell or in the same right cell. It is no harm to assume that y, w are in the same left cell since $\mu(y, w) = \mu(y^{-1}, w^{-1})$. Then y, w are in different right cell (cf [KL, §5]. This implies that $\gamma_{y^{-1},w,u} = 0$ for all $u \in \mathfrak{S}_n$ (see [L1, 6.3(a)]). If $\delta_{y^{-1},w,d} \neq 0$, then $d = d(w)$. Thus in this case we still have $\mu(y, w) = \delta_{y^{-1},w,d(w)}$.

Conversely, assume that $\delta_{x,y,z} \neq 0$. It is no harm to assume that $a(y) \geq a(x)$. Then y and z are in the same left cell, cf 1.2 (c) and [L1, 6.3(b)]. Thus we can find a sequence of right star operations $\varphi_k, \dots, \varphi_1$ such that $\varphi_k \dots \varphi_1(z) = d$ is in $\mathcal{D}_0$, see [KL, §5]. By a similar argument as for [X, Prop. 1.4.4(b)], we see that $h_{x,y,z} = h_{x,y',d}$, here $y' = \varphi_k \dots \varphi_1(y)$. In particular we have $\delta_{x,y,z} = \delta_{x,y',d}$. If $a(y) > a(x)$, by Springer's formula 1.2 (a), the value $\delta_{x,y',d}$ is equal to $\mu(x^{-1}, y')$ or $\mu(y', x^{-1})$. If $a(y) = a(x)$, then d and x are in the same right cell, d and y' are in the same left cell. So x, y'^{-1} are in the same right cell and are not in the same left cell. Thus $\gamma_{x,y',u} = 0$ for any u (see [L1, 6.3(a)]). In this case we also have $\delta_{x,y',d} = \mu(x^{-1}, y')$ or $\delta_{x,y',d} = \mu(y', x^{-1})$.

The proposition is proved.

In the proof we see the following

Corollary 2.2. If $a(y) \geq a(x)$, then $\delta_{x,y,z} = \delta_{x,y',d} = \mu(x^{-1}, y')$ or $\mu(y', x^{-1})$ for some y' in the right cell containing y and some d in $\mathcal{D}_0$.

When y and w are in the same two-sided cell we have the following result.

Proposition 2.3. The set $\{\pi(f)|f \in \mathcal{D}_1\}$ is the same as the set $\{\mu(x, y)|x \underset{LR}{\sim} y \text{ and } \mu(x, y) \neq 0\}$.

Proof: If $\mu(x, y) \neq 0$ and x and y are in the same two-sided cell, by 1.3 (b) and [L1, 6.3], x and y are in the same left cell or in the same right cell. It is no harm to assume that x and y are in the same left cell. By Springer's formula 1.2 (a), $\delta = \delta_{x, y^{-1}, d(y^{-1})} \neq 0$ or $\gamma = \gamma_{x, y^{-1}, f} \neq 0$ for some $f \in \mathcal{D}_1$. Since y and w are not in the same right cell, we have $\delta = 0$ and $\gamma \neq 0$. Note that $\gamma_{x, y^{-1}, f} \neq 0$ implies that $f \underset{R}{\sim} x$ and $f \underset{L}{\sim} y^{-1}$, see [L1, 6.3]. Such f is unique. Moreover, using [X, Theorem 1.4.5], [KL, §5] and [L2, 1.4(a)], $\gamma_{x, y^{-1}, f} = 1$ in this case. Therefore $\mu(x, y) = \gamma_{x, y^{-1}, f} \pi(f) = \pi(f)$. The proposition is proved.

For simplifying statements, in the rest of this section we set $\mu(y, w) = \mu(w, y)$ whenever one of $\mu(y, w)$, $\mu(w, y)$ is defined. Note that we have $\mu(f, d(f)) = \gamma_{f, d(f), f} \pi(f) = \pi(f)$. Thus we have

Corollary 2.4. The set $\{\mu(x, y)|x \underset{LR}{\sim} y \text{ and } \mu(x, y) \neq 0\}$ is the same as the set $\{\mu(f, d)|f \in \mathcal{D}_1, d \in \mathcal{D}_0, \text{ and } f \underset{LR}{\sim} d, \mu(f, d) \neq 0\}$.

2.5. Let $f, x \in \mathfrak{S}_n$. If $x \underset{LR}{\sim} f$ and $h_{f, f, x} \neq 0$, then $x \underset{L}{\sim} f$ and $x \underset{R}{\sim} f$. Thus we must have $x = f$, since in a two-sided cell of $\mathfrak{S}_n$, each left cell and each right cell have exactly one common element. Let M_f be the irreducible module of $H \otimes_A \mathbf{Q}(q^{\frac{1}{2}})$ provided by the left cell of $\mathfrak{S}_n$ containing f (cf, [KL]). Then we have

$$h_{f, f, f} = \text{tr}(C_f, M_f).$$

By [X, Prop. 1.4.5 (b)], we see that $h_{f, f, f} = h_{f, d, d}$, where $d \in \mathcal{D}_0$ and $d \underset{R}{\sim} f$. Thus $\delta_{f, f, f} = \mu(f, d)$. Therefore we also have $\{\mu(x, y)|x \underset{LR}{\sim} y\} = \{\delta_{f, f, f}|f \in \mathfrak{S}_n\}$ (we set $\mu(x, x) = 0$).

3. Proof of Theorem 1.4

3.1. Now we can prove the main result. Let $y \leq w$ be elements in $\mathfrak{S}_n$ such that $a(y) < a(w)$ and $\mu(y, w) \neq 0$.

By 1.3 (a), we have $y \underset{L}{\leq} w$ and $y \underset{R}{\leq} w$. Thus $L(y)$ is a subset of $L(w)$. If there is a simple reflection s in $L(w) - L(y)$, then $P_{y, w} = P_{sy, w}$ and the condition $\mu(y, w) \neq 0$ forces that $sy = w$ and $\mu(y, w) = 1$. Now assume that $L(w) = L(y)$. According to [KL, §5], there exists some subset I of $\{1, 2, \dots, n-1\}$ and a sequence $\varphi_i, \dots, \varphi_1$ of left star operations such that $\varphi_i \cdots \varphi_1(w) \underset{R}{\sim} w_I$. (We use W_I for the subgroup of $\mathfrak{S}_n$ generated by all s_i , $i \in I$, and denote by w_I the longest element of W_I .) For $j = 1, 2, \dots, i$, set $w_j = \varphi_j \cdots \varphi_1(w)$ and set $w_0 = w$.

Let k be the maximal number among $1, 2, \dots, i$ such that $y_j = \varphi_j \cdots \varphi_1(y)$ is well defined for all $j = 1, 2, \dots, k$. Set $y_0 = y$. If for some $1 \leq j \leq k$, we have $w_j \leq y_j$ and $y_m \leq w_m$ for $m = 0, 1, \dots, j-1$, using 1.3 (e) we see that $1 = \mu(w_j, y_j) = \mu(y_{j-1}, w_{j-1}) = \mu(y, w)$.

Now assume that $y_j \leq w_j$ for all $j = 1, 2, \dots, k$. Using 1.3 (e) we get $\mu(y_k, w_k) = \mu(y, w)$. We claim that $L(w_k) - L(y_k)$ is non-empty. Since $a(y_j) = a(y) < a(w) = a(w_j)$ for all $j = 0, 1, 2, \dots, k$, we always have $L(y_j) \subseteq L(w_j)$ for all j . If $k < i$, the set $L(w_k) - L(y_k)$ must be non-empty. Otherwise, we must have $L(w_k) = L(y_k)$. Thus $y_{k+1} = \varphi_{k+1}(y_k)$ is well defined. This contradicts the assumption on k . When $k = i$, we have $L(w_k) = \{s_j|j \in I\}$. If $L(w_k) \subseteq L(y_k)$, then we can find y' such that $y_k = w_I y'$ and $l(y_k) = l(w_I) + l(y')$. Thus $a(y) = a(y_k) \geq a(w_I) = a(w_k) = a(w)$. A

contradiction. Therefore in this case we also have that $L(w_k) - L(y_k)$ is non-empty. We then must have $l(w_k) = l(y_k) + 1$ and $\mu(y_k, w_k) = \mu(y, w) = 1$.

The theorem is proved.

Remark. Using the result [S, Lemma 18.3.2] and a similar argument as above one can prove that Theorem 1.4 remains true for an affine Weyl group of type $\tilde{A}_{n-1}$.

Proposition 3.2. For $\mathfrak{S}_n$, $n \geq 4$, the set $\{\mu(y, w) | y, w \in \mathfrak{S}_n\}$ is equal to the set $\{\mu(y, w) | y, w \in \mathfrak{S}_n \text{ and } a(y) > a(w)\}$.

Proof: Clearly, when $n \geq 4$, 1 is in the set $A = \{\mu(y, w) | y, w \in \mathfrak{S}_n \text{ and } a(y) > a(w)\}$. By Theorem 1.4, we only need to show that if $a(y) = a(w)$, then $\mu(y, w)$ is in A . To do this we need some preparation.

Let $U = U_0$ be the subgroup of $\mathfrak{S}_n$ generated by $s_2, s_3, \dots, s_{n-1}$. For $i = 1, 2, \dots, n-1$, we set $U_i = s_i \cdots s_2 s_1 U_0$ and $U'_i = U_0 s_1 s_2 \cdots s_i$. Then $U_i \cap U_j = \emptyset$ if $i \neq j$ and $\mathfrak{S}_n = U_0 \cup U_1 \cup \cdots \cup U_{n-1}$. Moreover if w is in U_0 , then the length of $s_i \cdots s_2 s_1 w$ is $l(w) + i$. The following assertions (a) and (b) are easy to prove.

(a) Let $y \in U_i$ and $w \in U_j$. If $y \leq w$, then $i \leq j$.

(b) Let $y = s_i \cdots s_2 s_1 y_1$ and $w = s_i \cdots s_2 s_1 w_1$ be elements in U_i . Then $P_{y,w} = P_{y_1, w_1}$.

For a subset I of $\{1, 2, \dots, n-1\}$, let W_I be the subgroup of $\mathfrak{S}_n$ generated by $s_i, i \in I$ and w_I the longest element of W_I .

Let $I = I_1 \cup I_2 \cup \cdots \cup I_m$ be a subset of $\{1, 2, \dots, n-1\}$. Assume that $I_1 = \{1, 2, \dots, k_1\}$ and $I_j = \{k_{j-1} + 2, k_{j-1} + 3, \dots, k_j\}$ for $j = 2, 3, \dots, m$. Then $w_I = w_{I_1} w_{I_2} \cdots w_{I_m}$. Assume that $k_1 + 1 \geq k_2 - k_1 \geq k_3 - k_2 \geq \cdots \geq k_m - k_{m-1} \geq 1$. The shape of the Young tableau $P(w_I)$ is $\lambda(w_I) = (k_1 + 1, k_2 - k_1, \dots, k_m - k_{m-1}, 1, \dots, 1)$.

According to 1.5 and [KL, §5], we have

(c) for each element w in $\mathfrak{S}_n$ we can find a unique subset I_λ of $\{1, 2, \dots, n-1\}$ as above such that $w \underset{LR}{\sim} w_I$ and can find a sequence $\varphi_i, \dots, \varphi_1$ of left star operations such that $\varphi_i \cdots \varphi_1(w) \underset{R}{\sim} w_I$.

Now we claim the following assertion.

(d) Let I be as above and $w \underset{R}{\sim} w_I$. Then w is in U_{k_1} .

We argue for (d). Let j be such that w is in U_j . Clearly we have $j \geq k_1$. If $j > k_1$, then $j = k_a$ for some $2 \leq a \leq m$ and $w = s_j s_{j-1} \cdots s_2 s_1 w_1$ for some $w_1 \in U$. Let $u = s_{k_{a-1}+2} s_{k_{a-1}+3} \cdots s_j w$. Then $w \underset{LR}{\leq} u$ and $s_{k_{a-1}+1} u \leq u$. Clearly, for any i in $I_1 \cup I_2 \cup \cdots \cup I_{a-1} \cup I_{a+1} \cup \cdots \cup I_m$, we have $s_i s_{k_{a-1}+2} s_{k_{a-1}+3} \cdots s_j = s_{k_{a-1}+2} s_{k_{a-1}+3} \cdots s_j s_i$. Thus, for these i we have $s_i u \leq u$. It is easy to check that for i in $I_a - \{k_{a-1} + 2\}$, we have $s_i u \leq u$. Let

$$J = I_1 \cup I_2 \cup \cdots \cup I_{a-2} \cup (I_{a-1} \cup \{k_{a-1} + 1\}) \cup (I_a - \{k_{a-1} + 2\}) \cup I_{a+1} \cup \cdots \cup I_m.$$

Then $u = w_J u_1$ for some u_1 with $l(u) = l(w_J) + l(u_1)$. Thus $a(u) \geq a(w_J) = a(w_I) + 1 + (k_{a-1} - k_{a-2}) - (k_a - k_{a-1}) \geq a(w_I) + 1$ (we set $k_{-1} = -1$). This contradicts $a(u) \leq a(w) = a(w_I)$. Therefore (d) is true.

Now we can complete the proof of the proposition. Assume that $a(y) = a(w)$. Then $y \underset{L}{\sim} w$ or $y \underset{R}{\sim} w$. It is no harm to assume that $y \underset{R}{\sim} w$ since $\mu(y, w) = \mu(y^{-1}, w^{-1})$. Also we may assume that $\mu(y, w) > 1$. Since a left star operation sends a right cell to a right cell with the same a -function value and it keeps the value of the $\mu(y, w)$ (see 1.3(e)), it is now harm to assume that $y \underset{R}{\sim} w \underset{R}{\sim} w_I$, where I is as in (d). By (d), we know that both y and w are in U_{k_1} . Set $y = s_{k_1} \cdots s_2 s_1 y_1$ and $w = s_{k_1} \cdots s_2 s_1 w_1$. Then y_1 and w_1 are in U . By (b), we have $\mu(y, w) = \mu(y_1, w_1)$.

If $a(y_1) < a(w_1)$ or $a(y_1) > a(w_1)$, we are done. If $a(y_1) = a(w_1)$, continuing the above process or using induction on n , we see that $\mu(y, w)$ is in A .

The proposition is proved.

Here is a question. For $y \leq w$ in $\mathfrak{S}_n$ with $\mu(y, w) \neq 0$ and $a(y) > a(w)$, whether we can find y' and w' in some $\mathfrak{S}_r$ such that $y' \underset{LR}{\sim} w'$ and $\mu(y, w) = \mu(y', w')$.

3.3. Using 1.5 (b) we get the following assertion.

(a) Let w be in $\mathfrak{S}_n$ and I be as (d) of the proof of Proposition 3.2. Then $w \underset{R}{\sim} w_I$ if and only if the following conditions are satisfied:

- (1) For $k_{i-1} + 2 \leq f < g \leq k_i$ (set $k_0 = -1$), we have $w^{-1}(f) > w^{-1}(g)$.
 - (2) For each $1 \leq f \leq n$, we have $|w(I_i) \cap \{1, 2, \dots, f\}| \geq |w(I_{i+1}) \cap \{1, 2, \dots, f\}|$. (We set $I_{m+j} = \{k_m + j\}$ for $1 \leq j \leq n - k_m$.)
- (b) By means of (b) in the proof of Proposition 3.2 we see easily that $\mu(y, w) \leq 1$ if $y \leq w$ and $a(w) = 1$.

Acknowledgement: Part of the work was done during my visit to the Department of Mathematics, Hong Kong University of Science and Technology. I am very grateful to the department for hospitality and for financial support. I would like to thank Professor Y. Zhu for helpful discussions and for arranging the visit.

References

- [BV] D. Barbasch and D. Vogan, *Primitive ideals and orbital integrals in complex classical groups*, Math. Ann. 259 (1982), 153-199.
- [GM] A. M. Garsia and T. J. McLarnan, *Relations between Young's Natural and the Kazhdan-Lusztig representations of S_n* , Adv. Math. 69 (1988), 32-92.
- [KL] D. Kazhdan and G. Lusztig, *Representations of Coxeter groups and Hecke algebras*, Invent. Math. 53 (1979), 165-184.
- [L1] G. Lusztig, *Cells in affine Weyl groups*, in "Algebraic groups and related topics", Advanced Studies in Pure Math., vol. 6, Kinokunia and North Holland, 1985, pp. 255-287.
- [L2] G. Lusztig, *Cells in affine Weyl groups, II*, J. Alg. 109 (1987), 536-548.
- [L3] G. Lusztig, *Letter to T. A. Springer*, 1987.
- [L4] G. Lusztig, *Nonlocal finiteness of a W -graph*, Represent. Theory 1 (1996), 25-30.
- [MW] T. McLarnan and G. Warrington, *Counterexamples to the 0,1-Conjecture*, Represent. Theory 7 (2003), 181-195.
- [S] J.-Y. Shi, *Kazhdan-Lusztig cells of certain affine Weyl groups*, LNM 1179, Springer-Verlag, Berlin, 1986.
- [Sp] T. A. Springer, *Letter to G. Lusztig*, 1987.
- [X] N. Xi, *The based ring of two-sided cells of affine Weyl groups of type $\tilde{A}_{n-1}$* , Mem. of AMS, No. 749, vol. 157, 2002.

N.X., INSTITUTE OF MATHEMATICS, CHINESE ACADEMY OF SCIENCES, BEIJING 100080, CHINA
E-mail address: nanhua@mail2.math.ac.cn